

## 2012 Appraisal Index

To assist clinicians looking for authoritative assistance with clinical problems, the journal publishes an annual index of content from the most recent two years of Appraisal pages. This index includes content from Volumes 57 and 58 of *Journal of Physiotherapy*. Content is indexed under the PEDro codes: subspecialty, intervention, problem, and body part, and identified by Appraisal section and Volume and page number. Some content is indexed under more than one code.

### SUBDISCIPLINE

#### Cardiothoracics

##### CAPs

Questioning the role of targeted respiratory physiotherapy over and above a standard clinical pathway in the postoperative management of patients following open thoracotomy	Vol 57 No 4, p 256
Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD)	Vol 58 No 1, p 60
Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease	Vol 58 No 2, p 129
Interval training confers greater gains than continuous training in people with heart failure	Vol 58 No 3, p 199
Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease	Vol 58 No 4, p 270
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271

#### Continence & Women's Health

##### CAPs

Manual lymph drainage when added to advice and exercise may not be effective in preventing lymphoedema after surgery for breast cancer	Vol 57 No 4, p 258
A 12-week exercise program performed during the second trimester does not prevent gestational diabetes in healthy pregnant women	Vol 58 No 3, p 198

##### Clinimetrics

2D real time ultrasound for pelvic floor muscle assessment	Vol 57 No 1, p 59
Active Straight Leg Raise	Vol 58 No 2, p 132

#### Gerontology

##### CAPs

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No4, p 269
--	-------------------

##### Clinical Practice Guidelines

Prevention of falls in older persons	Vol 57 No 2, p 130
Palliation in aged care	Vol 58 No 1, p 63

Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis	Vol 58 No 2, p 128
Osteoarthritis	Vol 58 No 2, p 133
Hip fracture	Vol 58 No 4, p 275
Osteoporosis	Vol 58 No 4, p 275

#### Musculoskeletal

##### CAPs

Telephone-based patient self-management program might be effective in reducing osteoarthritis-related pain	Vol 57 No 2, p 125
Lateral wedge insoles worn for 12 months provided no symptomatic or structural benefit for people with medial knee osteoarthritis	Vol 57 No 3, p 195
Surgery with disc prosthesis may produce better outcomes than multidisciplinary rehabilitation for patients with chronic low back pain	Vol 57 No 4, p 257
Targeted physiotherapy treatment for low back pain based on clinical risk can improve clinical and economic outcomes when compared with current best practice	Vol 58 No 1, p 57
A specific exercise program for patients with subacromial impingement syndrome can improve function and reduce the need for surgery	Vol 58 No2, p 127
Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis	Vol 58 No 2, p 128
Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training	Vol 58 No 3, p 200
Multi-modal realignment treatment decreases pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272

##### Clinimetrics

The Neer sign and Hawkins-Kennedy test for shoulder impingement	Vol 57 No 4, p 260
Western Ontario Rotator Cuff Index	Vol 58 No 3, p 201
The Patient-Rated Elbow Evaluation (PREE)	Vol 58 No 4, p 274

##### Clinical Practice Guidelines

Achilles pain, stiffness, and muscle power deficits	Vol 57 No 1, p 62
Knee pain	Vol 57 No 2, p 130
Arthroscopic shoulder surgery	Vol 57 No 4, p 261
Tension-type headache	Vol 57 No 4, p 261
Osteoarthritis	Vol 58 No 2, p 133
Hand, hip, and knee osteoarthritis	Vol 58 No 3, p 203
Cervical radiculopathy	Vol 58 No 3, p 203
Hip fracture	Vol 58 No 4, p 275
Osteoporosis	Vol 58 No 4, p 275

## Neurology

### CAPs

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia Vol 58 No 1, p 59

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility Vol 58 No4, p 269

### Clinimetrics

International standards for the neurological classification of spinal cord injury Vol 57 No 2, p 129

The Dix-Hallpike Test Vol 58 No 2, p 131

### Clinical Practice Guidelines

Early Parkinson's disease Vol 57 No 3, p 198

Tension-type headache Vol 57 No 4, p 260

Cervical radiculopathy Vol 58 No 3, p 203

## Orthopaedics

### CAPs

A specific exercise program for patients with subacromial impingement syndrome can improve function and reduce the need for surgery Vol 58 No2, p 127

Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis Vol 58 No 2, p 128

Multi-modal realignment treatment decreases pain in people with medial tibiofemoral osteoarthritis Vol 58 No 4, p 272

The Patient-Rated Elbow Evaluation (PREE) Vol 58 No 4, p 274

### Clinical Practice Guidelines

Osteoarthritis Vol 58 No 2, p 133

Hand, hip, and knee osteoarthritis Vol 58 No 3, p 203

Hip fracture Vol 58 No 4, p 275

Osteoporosis Vol 58 No 4, p 275

## Paediatrics

### CAPs

Aerobic exercise enhances executive function and academic achievement in sedentary, overweight children aged 7–11 years Vol 57 No 4, p 255

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia Vol 58 No 1, p 59

Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy Vol 58 No 3, p 197

### Clinimetrics

The six-minute walk test in paediatric populations Vol 57 No 2, p128

### Clinical Practice Guidelines

Developmental co-ordination disorder Vol 58 No 2, p 133

## Sports

### CAPs

Eccentric hamstring muscle training can prevent hamstring injuries in soccer players Vol 58 No 1, p 58

## Other

### Clinimetrics

Pain Intensity Ratings Vol 58 No 1, p 61

The Dix-Hallpike Test Vol 58 No 2, p 131

Brief Illness Perception Questionnaire (Brief IPQ) Vol 58 No 3, p 202

SPHERE 12 Screening Questionnaire Vol 58 No 4, p 273

# INTERVENTION

## Acupuncture

### CAPs

Traditional Chinese Acupuncture was not superior to sham acupuncture for knee osteoarthritis but delivering treatment with high expectations of improvement was superior to delivering treatment with neutral expectations Vol 57 No 1, p 56

## Behaviour Modification

### CAPs

Telephone-based patient self-management program might be effective in reducing osteoarthritis-related pain Vol 57 No 2, p 125

## Fitness Training

### CAPs

Supervised aerobic and resistance exercise improves glycaemic control and modifiable cardiovascular risk factors in people with Type 2 diabetes mellitus Vol 57 No 2, p 126

Aerobic exercise enhances executive function and academic achievement in sedentary, overweight children aged 7–11 years Vol 57 No 4, p 255

Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease Vol 58 No 2, p 129

Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy Vol 58 No 3, p 197

A 12-week exercise program performed during the second trimester does not prevent gestational diabetes in healthy pregnant women Vol 58 No 3, p 198

Interval training confers greater gains than continuous training in people with heart failure Vol 58 No 3, p 199

Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training Vol 58 No 3, p 200

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271
Multi-modal realignment treatment decreases pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272

## Orthoses, Taping, Splinting

### CAPs

Foot orthoses can reduce lower limb overuse injury rate	Vol 57 No 3, p 193
Lateral wedge insoles worn for 12 months provided no symptomatic or structural benefit for people with medial knee osteoarthritis	Vol 57 No 3, p 195

## Respiratory Therapy

### CAPs

Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD)	Vol 58 No 1, p 60
Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease	Vol 58 No 4, p 270

## Strength Training

### CAPs

Eccentric hamstring muscle training can prevent hamstring injuries in soccer players	Vol 58 No 1, p 58
Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease	Vol 58 No 2, p 129
Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy	Vol 58 No 3, p 197
Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training	Vol 58 No 3, p 200

## Stretching, Mobilisation, Manipulation, Massage

### CAPs

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271
Multi-modal realignment treatment decreases pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272

## PROBLEM

### Difficulty with Sputum Clearance

#### Clinical Practice Guidelines

Bronchiectasis	Vol 57 No 3, p 198
----------------	--------------------

### Impaired Ventilation

#### CAPs

Using titrated oxygen instead of high flow oxygen during an acute exacerbation of chronic obstructive pulmonary disease (COPD) saves lives	Vol 57 No 1, p 55
Patients with chronic obstructive pulmonary disease (COPD) who are not hypoxaemic at rest do not benefit from home oxygen	Vol 57 No 2, p 127
Resistance training preserves skeletal muscle function in patients with COPD who are hospitalised with an acute exacerbation	Vol 57 No 3, p 194
Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD)	Vol 58 No 1, p 60
Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease	Vol 58 No 4, p 270

#### Clinical Practice Guidelines

Bronchiectasis	Vol 57 No 3, p 198
----------------	--------------------

### Incontinence

#### Clinimetrics

2D real time ultrasound for pelvic floor muscle assessment	Vol 57 No 1, p 59
--	-------------------

### Muscle Weakness

#### CAPs

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia	Vol 58 No 1, p 59
Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy	Vol 58 No 3, p 197
Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training	Vol 58 No 3, p 200
Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271
Multi-modal realignment treatment decreases pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272

**Clinimetrics**

Rehabilitative ultrasound imaging	Vol 57 No 3, p 196
Western Ontario Rotator Cuff Index	Vol 58 No 3, p 201

**Pain****CAPs**

Telephone-based patient self-management program might be effective in reducing osteoarthritis-related pain	Vol 57 No 2, p 125
Multi-modal realignment treatment decreases pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272

**Clinimetrics**

(Thermal) Quantitative Sensory Testing–tQST	Vol 57 No 1, p 58
Pain Intensity Ratings	Vol 58 No 1, p 61
Active Straight Leg Raise	Vol 58 No 2, p 132
Western Ontario Rotator Cuff Index	Vol 58 No 3, p 201
The Patient-Rated Elbow Evaluation (PREE)	Vol 58 No 4, p 274

**Clinical Practice Guidelines**

Osteoarthritis	Vol 58 No 2, p 133
Hand, hip, and knee osteoarthritis	Vol 58 No 3, p 203
Cervical radiculopathy	Vol 58 No 3, p 203

**Reduced Exercise Tolerance****CAPs**

Interval training confers greater gains than continuous training in people with heart failure	Vol 58 No 3, p 199
Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271

**Clinimetrics**

The six-minute walk test in paediatric populations	Vol 57 No 2, p 128
Rating of Perceived Exertion (RPE)	Vol 58 No 1, p 62

**Other****CAPs**

Targeted physiotherapy treatment for low back pain based on clinical risk can improve clinical and economic outcomes when compared with current best practice	Vol 58 No 1, p 57
Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis	Vol 58 No 2, p 128
Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease	Vol 58 No 2, p 129
Exercise training improves cardiovascular fitness in people receiving haemodialysis for chronic renal disease	Vol 58 No 2, p 130

**Clinimetrics**

Rating of Perceived Exertion (RPE)	Vol 58 No 1, p 62
The Dix-Hallpike Test	Vol 58 No 2, p 131
Brief Illness Perception Questionnaire (Brief IPQ)	Vol 58 No 3, p 202
SPHERE 12 Screening Questionnaire	Vol 58 No 4, p zz

**Clinical Practice Guidelines**

Palliation in aged care	Vol 58 No 1, p 63
Hip fracture	Vol 58 No 4, p 275
Osteoporosis	Vol 58 No 4, p 275

**BODY PART****Head & Neck****Clinimetrics**

The Dix-Hallpike Test	Vol 58 No 2, p 131
-----------------------	--------------------

**Upper Arm, Shoulder or Shoulder Girdle****Clinimetrics**

Shoulder Pain and Disability Index (SPADI)	Vol 57 No 3, p 197
The Neer sign and Hawkins-Kennedy test for shoulder impingement	Vol 57 No 4, p 260
Western Ontario Rotator Cuff Index	Vol 58 No 3, p 201
The Patient-Rated Elbow Evaluation (PREE)	Vol 58 No 4, p 274

**Clinical Practice Guidelines**

Arthroscopic shoulder surgery	Vol 57 No 4, p 261
A specific exercise program for patients with subacromial impingement syndrome can improve function and reduce the need for surgery	Vol 58 No 2, p 127

**Hand or Wrist****CAPs**

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia	Vol 58 No 1, p 59
--	-------------------

**Clinical Practice Guidelines**

Hand, hip, and knee osteoarthritis	Vol 58 No 3, p 203
------------------------------------	--------------------

**Chest****CAPs**

Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD)	Vol 58 No 1, p 60
Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease	Vol 58 No 2, p 129
Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease	Vol 58 No 4, p 270



## Thoracic Spine

### Clinimetrics

- International standards for the neurological classification of spinal cord injury Vol 57 No 2, p 129

### Clinical Practice Guidelines

- Cervical radiculopathy Vol 58 No 3, p 203

## Lumbar Spine, SIJ or Pelvis

### CAPs

- Surgery with disc prosthesis may produce better outcomes than multidisciplinary rehabilitation for patients with chronic low back pain Vol 57 No 4, p 257
- Targeted physiotherapy treatment for low back pain based on clinical risk can improve clinical and economic outcomes when compared with current best practice Vol 58 No 1, p 57
- Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training Vol 58 No 3, p 200

### Clinimetrics

- International standards for the neurological classification of spinal cord injury Vol 57 No 2, p 129
- Active Straight Leg Raise Vol 58 No 2, p 132

## Perineum or Genito-Urinary System

### Clinimetrics

- 2D real time ultrasound for pelvic floor muscle assessment Vol 57 No 1, p 59
- Exercise training improves cardiovascular fitness in people receiving haemodialysis for chronic renal disease Vol 58 No 2, p 130

## Thigh or Hip

### CAPs

- Eccentric hamstring muscle training can prevent hamstring injuries in soccer players Vol 58 No 1, p 58
- Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis Vol 58 No 4, p 272

### Clinical Practice Guidelines

- Hand, hip, and knee osteoarthritis Vol 58 No 3, p 203
- Hip fracture Vol 58 No 4, p 275

## Lower Leg or Knee

### CAPs

- Achilles pain, stiffness, and muscle power deficits Vol 57 No 1, p 62

- Lateral wedge insoles worn for 12 months provided no symptomatic or structural benefit for people with medial knee osteoarthritis Vol 57 No 3, p 195

- Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis Vol 58 No 2, p 128

### Clinical Practice Guidelines

- Knee pain Vol 57 No 2, p 130
- Osteoarthritis Vol 58 No 2, p 133
- Hand, hip, and knee osteoarthritis Vol 58 No 3, p 203

## Foot or Ankle

### CAPs

- Achilles pain, stiffness, and muscle power deficits Vol 57 No 1, p 62
- Foot orthoses can reduce lower limb overuse injury rate Vol 57 No 3, p 193
- Lateral wedge insoles worn for 12 months provided no symptomatic or structural benefit for people with medial knee osteoarthritis Vol 57 No 3, p 195

## Whole Body/Other

### CAPs

- Telephone-based patient self-management program might be effective in reducing osteoarthritis-related pain Vol 57 No 2, p 125
- Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy Vol 58 No 3, p 197
- A 12-week exercise program performed during the second trimester does not prevent gestational diabetes in healthy pregnant women Vol 58 No 3, p 198
- Interval training confers greater gains than continuous training in people with heart failure Vol 58 No 3, p 199
- Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke Vol 58 No 4, p 271

### Clinimetrics

- Brief Illness Perception Questionnaire (Brief IPQ) Vol 58 No 3, p 202
- SPHERE 12 Screening Questionnaire Vol 58 No 4, p 273

### Clinical Practice Guidelines

- General Health Questionnaire – 28 (GHQ-28) Vol 57 No 4, p 259
- Palliation in aged care Vol 58 No 1, p 63
- Type 2 diabetes Vol 58 No 1, p 63
- Developmental co-ordination disorder Vol 58 No 2, p 133
- Osteoporosis Vol 58 No 4, p 275